

Texture 4 zero Fritzsch-like lepton mass matrices

Gulsheen Ahuja¹, Sanjeev Kumar², Monika Randhawa³,
Manmohan Gupta^{1,*}, S. Dev²

¹ *Department of Physics, Centre of Advanced Study, P.U., Chandigarh, India.*

² *Department of Physics, H.P.U., Shimla, India.*

³ *University Institute of Engineering and Technology, P.U., Chandigarh, India.*

February 21, 2008

Abstract

For Majorana or Dirac neutrinos, using Fritzsch-like texture 4 zero mass matrices with parallel texture structures for the charged leptons and the Dirac neutrino mass matrix ($M_{\nu D}$), detailed predictions for cases pertaining to normal/inverted hierarchy as well as degenerate scenarios of neutrino masses have been carried out. The inverted hierarchy as well as degenerate scenarios seem to be ruled out at 3σ C.L. for both Majorana and Dirac neutrinos. For normal hierarchy, Jarlskog's rephasing invariant parameter J , the CP violating Dirac-like phase δ and the effective neutrino mass $\langle m_{ee} \rangle$ have been calculated. For this case, lower limits of m_{ν_1} and θ_{13} would have implications for the nature of neutrinos.

In the last few years, impressive advances have been made in understanding the phenomenology of neutrino oscillations through solar neutrino experiments [1], atmospheric neutrino experiments [2], reactor based experiments [3] and accelerator based experiments [4]. At present, one of the key issues in the context of neutrino oscillation phenomenology is to understand the pattern of neutrino masses and mixings which seems to be vastly different from that of quark masses and mixings. In the case of quarks the masses and mixing angles show distinct hierarchy, whereas in the case of neutrinos neither the mixing angles nor the neutrino masses show any distinct hierarchy. In fact, the two mixing angles governing solar and atmospheric neutrino oscillations look to be rather large, the third angle may be very small compared to these. Further, at present there is no consensus about neutrino masses which may show normal/inverted hierarchy or may even be degenerate. Furthermore, the situation becomes complicated when one realizes that neutrino masses are much smaller than charged lepton and quark masses.

*mmgupta@pu.ac.in

Taking clue from the success of texture specific mass matrices in the context of quarks [5], several attempts [6, 7] have been made to consider similar lepton mass matrices for explaining the pattern of neutrino masses and mixings. In the absence of sufficient amount of data regarding neutrino masses and mixing angles, it would require a very careful scrutiny of all possible textures to find viable structures which are compatible with data and theoretical ideas so that these be kept in mind while formulating mass matrices at the GUT (Grand Unified Theories) scale. In this context, most of the attempts to understand the pattern of neutrino masses and mixings have been carried out using the seesaw mechanism [8] given by

$$M_\nu = -M_{\nu D}^T (M_R)^{-1} M_{\nu D}, \quad (1)$$

where $M_{\nu D}$ and M_R are respectively the Dirac neutrino mass matrix and the right-handed Majorana neutrino mass matrix. In this context, it should be noted that the predictions are somewhat different, on the one hand when texture is imposed only on $M_{\nu D}$ and M_R and on the other hand when M_ν and $M_{\nu D}$ have the same texture by imposing ‘texture invariant conditions’ [9, 10]. Similarly, as compared to the Majorana case, the predictions are different for Dirac neutrinos even if $M_{\nu D}$ has the same texture as that of M_ν .

Texture 4 zero mass matrices are known to explain the pattern of quark masses and mixings [11] as well as are known to be compatible with specific models of GUTs [5, 6] and these could be obtained using considerations of Abelian family symmetries [12]. Assuming normal hierarchy of masses as well as imposing texture 4 zero structure on $M_{\nu D}$ and charged lepton mass matrices, Xing *et al.* [9] have not only shown the compatibility of these with neutrino oscillation phenomenology but have also shown the seesaw invariance of these structures under certain conditions. Very recently, Matsuda *et al.* [10] have reiterated the success of texture 4 zero mass matrices in the case of quarks by showing that these mass matrices can accommodate the present value of $\sin 2\beta$. Also, for normal hierarchy they have shown that texture 4 zero lepton mass matrices can accommodate large values of mixing angle s_{13} . In particular, by imposing texture invariant conditions they have shown that M_ν can be texture 2 zero when one assumes Fritzsch-like texture 2 zero structure for $M_{\nu D}$, M_R as well as for charged lepton mass matrix.

In view of the success of texture 4 zero mass matrices in the context of quark mixing phenomenology, it would be interesting to carry out an exhaustive and detailed analysis of these in the case of leptons. In particular, for both Majorana and Dirac neutrinos it would be interesting to investigate the compatibility of texture 4 zero lepton mass matrices with the inverted hierarchy and degenerate scenario of neutrino masses. In the case of Majorana neutrinos due to the absence of any guidelines for M_R , to keep the number of independent parameters under control, it would perhaps be desirable to keep its structure as simple as possible. It would also be desirable to study the implications of these mass matrices when texture is imposed only on $M_{\nu D}$ and not on M_ν .

Considering Fritzsch-like texture 4 zero lepton mass matrices, with neutrinos and charged leptons having parallel texture structures, the purpose of the present communication is to investigate in detail the implications of these for normal/inverted hierarchy as

well as ‘different’ degenerate scenarios of neutrino masses. In particular, for the inverted hierarchy and degenerate scenarios, the implications of these structures have been carried out for both Majorana and Dirac neutrinos by exploring the parameter space available to any of the two mixing angles found by giving full variation to other parameters and phases. In the case of normal hierarchy, we have calculated several phenomenological quantities such as Jarlskog’s rephasing invariant parameter J , the CP violating Dirac-like phase δ and the effective neutrino mass $\langle m_{ee} \rangle$, related to neutrinoless double beta decay $(\beta\beta)_{0\nu}$.

To begin with, we define the modified Fritzsch-like matrices, e.g.,

$$M_l = \begin{pmatrix} 0 & A_l & 0 \\ A_l^* & D_l & B_l \\ 0 & B_l^* & C_l \end{pmatrix}, \quad M_{\nu D} = \begin{pmatrix} 0 & A_\nu & 0 \\ A_\nu^* & D_\nu & B_\nu \\ 0 & B_\nu^* & C_\nu \end{pmatrix}, \quad (2)$$

M_l and $M_{\nu D}$ respectively corresponding to Dirac-like charged lepton and neutrino mass matrices. The above matrices are texture 4 zero type with D_l and D_ν being non-zero along with $A_{l(\nu)} = |A_{l(\nu)}|e^{i\alpha_{l(\nu)}}$ and $B_{l(\nu)} = |B_{l(\nu)}|e^{i\beta_{l(\nu)}}$, in case these are symmetric then $A_{l(\nu)}^*$ and $B_{l(\nu)}^*$ should be replaced by $A_{l(\nu)}$ and $B_{l(\nu)}$, as well as $C_{l(\nu)}$ and $D_{l(\nu)}$ should respectively be defined as $C_{l(\nu)} = |C_{l(\nu)}|e^{i\gamma_{l(\nu)}}$ and $D_{l(\nu)} = |D_{l(\nu)}|e^{i\omega_{l(\nu)}}$.

To fix the notations and conventions as well as to facilitate the understanding of inverted hierarchy case and its relationship to the normal hierarchy case, we detail the formalism connecting the mass matrix to the neutrino mixing matrix. The mass matrices M_l and $M_{\nu D}$ given in equation (2), for Hermitian as well as symmetric case, can be exactly diagonalized. Details of Hermitian case can be looked up in our earlier work [13], the symmetric case can similarly be worked out. To facilitate diagonalization, the mass matrix M_k , where $k = l, \nu D$, can be expressed as

$$M_k = Q_k M_k^r P_k \quad (3)$$

or

$$M_k^r = Q_k^\dagger M_k P_k^\dagger, \quad (4)$$

where M_k^r is a real symmetric matrix with real eigenvalues and Q_k and P_k are diagonal phase matrices. For the hermitian case $Q_k = P_k^\dagger$, whereas for the symmetric case under certain conditions $Q_k = P_k$. In general, the real matrix M_k^r is diagonalized by the orthogonal transformation O_k , e.g.,

$$M_k^{diag} = O_k^T M_k^r O_k, \quad (5)$$

which on using equation (4) can be rewritten as

$$M_k^{diag} = O_k^T Q_k^\dagger M_k P_k^\dagger O_k. \quad (6)$$

To facilitate the construction of diagonalization transformations for different hierarchies, we introduce a diagonal phase matrix ξ_k defined as $\text{diag}(1, e^{i\pi}, 1)$ for the case of normal

hierarchy and as $\text{diag}(1, e^{i\pi}, e^{i\pi})$ for the case of inverted hierarchy. Equation (6) can now be written as

$$\xi_k M_k^{diag} = O_k^T Q_k^\dagger M_k P_k^\dagger O_k, \quad (7)$$

which can also be expressed as

$$M_k^{diag} = \xi_k^\dagger O_k^T Q_k^\dagger M_k P_k^\dagger O_k. \quad (8)$$

Making use of the fact that $O_k^* = O_k$ it can be further expressed as

$$M_k^{diag} = (Q_k O_k \xi_k)^\dagger M_k (P_k^\dagger O_k), \quad (9)$$

from which one gets

$$M_k = Q_k O_k \xi_k M_k^{diag} O_k^T P_k. \quad (10)$$

The case of leptons is fairly straight forward, for the neutrinos the diagonalizing transformation is hierarchy specific as well as requires some fine tuning of the phases of the right handed neutrino mass matrix M_R . To clarify this point further, in analogy with equation (10), we can express $M_{\nu D}$ as

$$M_{\nu D} = Q_{\nu D} O_{\nu D} \xi_{\nu D} M_{\nu D}^{diag} O_{\nu D}^T P_{\nu D}. \quad (11)$$

Substituting the above value of $M_{\nu D}$ in equation (1) one obtains

$$M_\nu = -(Q_{\nu D} O_{\nu D} \xi_{\nu D} M_{\nu D}^{diag} O_{\nu D}^T P_{\nu D})^T (M_R)^{-1} (Q_{\nu D} O_{\nu D} \xi_{\nu D} M_{\nu D}^{diag} O_{\nu D}^T P_{\nu D}). \quad (12)$$

On using $P_{\nu D}^T = P_{\nu D}$, the above equation can further be written as

$$M_\nu = -P_{\nu D} O_{\nu D} M_{\nu D}^{diag} \xi_{\nu D} O_{\nu D}^T Q_{\nu D}^T (M_R)^{-1} Q_{\nu D} O_{\nu D} \xi_{\nu D} M_{\nu D}^{diag} O_{\nu D}^T P_{\nu D}. \quad (13)$$

Assuming fine tuning, the phase matrices $Q_{\nu D}^T$ and $Q_{\nu D}$ along with $-M_R$ can be taken as $m_R \text{diag}(1, 1, 1)$ as well as using the unitarity of $\xi_{\nu D}$ and orthogonality of $O_{\nu D}$, the above equation can be expressed as

$$M_\nu = P_{\nu D} O_{\nu D} \frac{(M_{\nu D}^{diag})^2}{(m_R)^{-1}} O_{\nu D}^T P_{\nu D}. \quad (14)$$

The lepton mixing matrix, obtained from the matrices used for diagonalizing the mass matrices M_l and M_ν , is expressed as

$$U = (Q_l O_l \xi_l)^\dagger (P_{\nu D} O_{\nu D}). \quad (15)$$

Eliminating the phase matrix ξ_l by redefinition of the charged lepton phases, the above equation becomes

$$U = O_l^\dagger Q_l P_{\nu D} O_{\nu D}, \quad (16)$$

where $Q_l P_{\nu D}$, without loss of generality, can be taken as $(e^{i\phi_1}, 1, e^{i\phi_2})$, ϕ_1 and ϕ_2 being

related to the phases of mass matrices and can be treated as free parameters.

To understand the relationship between diagonalizing transformations for different hierarchies of neutrino masses as well as their relationship with the charged lepton case, we first consider the general diagonalizing transformation O_k , whose elements can be written as

$$\begin{aligned}
O_k(11) &= \sqrt{\frac{m_2 m_3 (m_3 + m_2 - D_{l(\nu)})}{(m_1 + m_2 + m_3 - D_{l(\nu)})(m_1 - m_3)(m_1 - m_2)}} \\
O_k(12) &= \sqrt{\frac{m_1 m_3 (m_1 + m_3 - D_{l(\nu)})}{(m_1 + m_2 + m_3 - D_{l(\nu)})(m_2 - m_3)(m_2 - m_1)}} \\
O_k(13) &= \sqrt{\frac{m_1 m_2 (m_2 + m_1 - D_{l(\nu)})}{(m_1 + m_2 + m_3 - D_{l(\nu)})(m_3 - m_2)(m_3 - m_1)}} \\
O_k(21) &= \sqrt{\frac{m_1 (m_3 + m_2 - D_{l(\nu)})}{(m_3 - m_1)(m_1 - m_2)}} \\
O_k(22) &= \sqrt{\frac{m_2 (m_3 + m_1 - D_{l(\nu)})}{(m_2 - m_3)(m_2 - m_1)}} \\
O_k(23) &= \sqrt{\frac{m_3 (m_2 + m_1 - D_{l(\nu)})}{(m_2 - m_3)(m_3 - m_1)}} \\
O_k(31) &= \sqrt{\frac{m_1 (m_1 + m_2 - D_{l(\nu)})(m_1 - m_3)(m_1 + m_3 - D_{l(\nu)})}{(m_1 - m_3)^2 (m_1 - m_2)(m_1 + m_2 + m_3 - D_{l(\nu)})}} \\
O_k(32) &= \sqrt{\frac{m_2 (D_{l(\nu)} - m_1 - m_2)(D_{l(\nu)} - m_2 - m_3)}{(m_1 + m_2 + m_3 - D_{l(\nu)})(m_2 - m_3)(m_2 - m_1)}} \\
O_k(33) &= \sqrt{\frac{m_3 (D_{l(\nu)} - m_2 - m_3)(D_{l(\nu)} - m_1 - m_3)}{(m_1 + m_2 + m_3 - D_{l(\nu)})(m_3 - m_1)(m_3 - m_2)}}, \tag{17}
\end{aligned}$$

where m_1, m_2, m_3 are eigenvalues of M_k . In the case of charged leptons, because of the hierarchy $m_e \ll m_\mu \ll m_\tau$, the mass eigenstates can be approximated respectively to the flavor eigenstates as has been considered by several authors [6, 14]. Using the approximation, $m_{l1} \simeq m_e$, $m_{l2} \simeq m_\mu$ and $m_{l3} \simeq m_\tau$, the first element of the matrix O_l can be obtained from the corresponding element of equation (17) by replacing m_1, m_2, m_3 by $m_e, -m_\mu, m_\tau$, e.g.,

$$O_l(11) = \sqrt{\frac{m_\mu m_\tau (m_\tau - m_\mu - D_l)}{(m_e - m_\mu + m_\tau - D_l)(m_\tau - m_e)(m_e + m_\mu)}}. \tag{18}$$

For normal hierarchy defined as $m_{\nu_1} < m_{\nu_2} \ll m_{\nu_3}$, as well as for the corresponding degenerate case given by $m_{\nu_1} \lesssim m_{\nu_2} \sim m_{\nu_3}$, equation (17) can also be used to obtain the

first element of diagonalizing transformation for Majorana neutrinos. By replacing m_1, m_2, m_3 by $\sqrt{m_{\nu 1} m_R}, -\sqrt{m_{\nu 2} m_R}, \sqrt{m_{\nu 3} m_R}$ in the equation, we get

$$O_\nu(11) = \sqrt{\frac{\sqrt{m_{\nu 2}} \sqrt{m_{\nu 3}} (\sqrt{m_{\nu 3}} - \sqrt{m_{\nu 2}} - D_\nu)}{(\sqrt{m_{\nu 1}} - \sqrt{m_{\nu 2}} + \sqrt{m_{\nu 3}} - D_\nu)(\sqrt{m_{\nu 3}} - \sqrt{m_{\nu 1}})(\sqrt{m_{\nu 1}} + \sqrt{m_{\nu 2}})}}, \quad (19)$$

where $m_{\nu 1}, m_{\nu 2}$ and $m_{\nu 3}$ are neutrino masses. The parameter D_ν is to be divided by $\sqrt{m_R}$, however as D_ν is arbitrary therefore we retain it as it is.

In the same manner, one can obtain the elements of diagonalizing transformation for the inverted hierarchy case defined as $m_{\nu 3} \ll m_{\nu 1} < m_{\nu 2}$ as well as for the corresponding degenerate case given by $m_{\nu 3} \sim m_{\nu 1} \lesssim m_{\nu 2}$. By replacing m_1, m_2, m_3 in equation (17) with $\sqrt{m_{\nu 1} m_R}, -\sqrt{m_{\nu 2} m_R}, -\sqrt{m_{\nu 3} m_R}$, we obtain

$$O_\nu(11) = \sqrt{\frac{\sqrt{m_{\nu 2}} \sqrt{m_{\nu 3}} (D_\nu + \sqrt{m_{\nu 2}} + \sqrt{m_{\nu 3}})}{(-\sqrt{m_{\nu 1}} + \sqrt{m_{\nu 2}} + \sqrt{m_{\nu 3}} + D_\nu)(\sqrt{m_{\nu 1}} + \sqrt{m_{\nu 3}})(\sqrt{m_{\nu 1}} + \sqrt{m_{\nu 2}})}}. \quad (20)$$

The other elements of diagonalizing transformations in the case of neutrinos as well as charged leptons can similarly be found. The above formalism has been presented for Majorana neutrinos, for the Dirac case the mixing matrix can easily be derived from diagonalizing transformation of M_l and $M_{\nu D}$.

It may be of interest to mention that in the case of normal hierarchy, the formulation of Matsuda *et al.* [10] can easily be obtained from the present case. For example, their element O_{11} can be obtained from our $O_k(11)$ by replacing ' d_f ' by ' $m_1 + m_2 + m_3 + D_{l(\nu)}$ ', similarly their other elements can be derived from the elements of our O_k . Further, it should also be noted that they have treated the 3×3 element of the mass matrix as free parameter whereas in the present case we have treated 2×2 element as a free parameter. Furthermore, we have not put any conditions so as to obtain a particular texture for M_ν .

Before discussing the results, we would like to mention some of the details pertaining to various inputs. For the purpose of our analysis we have used the results obtained by a recent global analysis carried out by Valle [15], incorporating solar [1], atmospheric [2], reactor [3] and accelerator based experiments [4]. The 3σ values of the neutrino mass and mixing parameters so obtained are

$$\Delta m_{12}^2 = (7.1 - 8.9) \times 10^{-5} \text{ eV}^2, \quad \Delta m_{23}^2 = (2.0 - 3.2) \times 10^{-3} \text{ eV}^2, \quad (21)$$

$$\sin^2 \theta_{12} = 0.24 - 0.40, \quad \sin^2 \theta_{23} = 0.34 - 0.68, \quad \sin^2 \theta_{13} \leq 0.040. \quad (22)$$

These values are quite compatible with those obtained very recently by a global analysis carried out by Garcia and Maltoni [16]. It may be mentioned that the present upper bound on $\sin \theta_{13}$ is somewhat lower than the CHOOZ bound [17]. Further, for the purpose of calculations, we have taken the lightest neutrino mass, the phases ϕ_1, ϕ_2 and $D_{l,\nu}$ as free parameters, the other two masses are constrained by $\Delta m_{12}^2 = m_{\nu 2}^2 - m_{\nu 1}^2$ and $\Delta m_{23}^2 = m_{\nu 3}^2 - m_{\nu 2}^2$ in the normal hierarchy case and by $\Delta m_{23}^2 = m_{\nu 2}^2 - m_{\nu 3}^2$ in the

inverted hierarchy case. It may be noted that lightest neutrino mass corresponds to m_{ν_1} for the normal hierarchy case and to m_{ν_3} for the inverted hierarchy case. For all the three hierarchies, the explored range of the lightest neutrino mass is taken to be $10^{-8} \text{ eV} - 10^{-1} \text{ eV}$, our conclusions remain unaffected even if the range is extended further. In the absence of any constraint on the phases, ϕ_1 and ϕ_2 have been given full variation from 0 to 2π . Although $D_{l,\nu}$ are free parameters, however, they have been constrained such that diagonalizing transformations O_l and O_ν always remain real. This implies, for leptons $-(m_{l_2} - m_{l_1}) < D_l < (m_{l_3} + m_{l_2})$, for Dirac neutrinos $-(m_{\nu_2} - m_{\nu_1}) < D_\nu < (m_{\nu_3} - m_{\nu_2})$ for normal hierarchy and $-(m_{\nu_2} - m_{\nu_1}) < D_\nu < (m_{\nu_1} - m_{\nu_3})$ for inverted hierarchy. Similarly, for Majorana neutrinos $-(\sqrt{m_{\nu_2}} - \sqrt{m_{\nu_1}}) < D_\nu < (\sqrt{m_{\nu_3}} - \sqrt{m_{\nu_2}})$ for normal hierarchy and $-(\sqrt{m_{\nu_2}} - \sqrt{m_{\nu_1}}) < D_\nu < (\sqrt{m_{\nu_1}} - \sqrt{m_{\nu_3}})$ for inverted hierarchy. The calculations pertaining to the case when charged leptons are in flavor basis can easily be deduced from the above calculations for both Majorana and Dirac neutrinos.

Considering Majorana or Dirac neutrinos, we have carried out detailed calculations pertaining to texture 4 zero lepton mass matrices for the possibilities of neutrino masses having normal/inverted hierarchy or being degenerate. To begin with, we consider the inverted hierarchy case for both types of neutrinos. In this context, it may be mentioned that for both the possibilities texture is imposed only on $M_{\nu D}$, with no such restriction on M_ν for the Majorana case. In figures 1a, 1b and 1c, for Majorana neutrinos we have plotted the parameter space corresponding to any of the two mixing angles by constraining the third angle by its values given in equation (22) while giving full allowed variation to other parameters. Also included in the figures are blank rectangular regions indicating the experimentally allowed 3σ region of the plotted angles. Interestingly, a general look at these figures reveals that the case of inverted hierarchy seems to be ruled out. A closer look at these figures brings out several interesting points. From figure 1a showing the plot of angles θ_{12} versus θ_{23} , one can immediately conclude that the plotted parameter space includes the experimentally allowed range of $\theta_{23} = 35.7^\circ - 55.6^\circ$, however it excludes the experimentally allowed range of $\theta_{12} = 29.3^\circ - 39.25^\circ$. This clearly indicates that at 3σ C.L. inverted hierarchy is not viable. It may be noted that while plotting this figure, θ_{13} is restricted by the bound given in equation (22), while Δm_{12}^2 and Δm_{23}^2 are constrained by the experimental limits given in equation (21). It may also be mentioned that although the 3σ upper limit of angle θ_{12} is not included in the plotted parameter space, yet it lies very near to the boundary, therefore the above conclusions needs to be checked further.

The conclusions arrived above can be further checked from figures 1b and 1c wherein we have plotted θ_{12} versus θ_{13} and θ_{23} versus θ_{13} respectively by constraining angles θ_{23} and θ_{12} . Both the figures indicate that the plotted parameter space does not include simultaneously the experimental bounds of the plotted angles, e.g., θ_{12} in the case of figure 1b and θ_{13} in figure 1c. Here it needs to be mentioned that similar to figure 1a, in figure 1b also the 3σ upper limit of angle θ_{12} lies very near to the boundary of the plotted parameter space, however in figure 1c the 3σ upper limit of angle θ_{13} is well below the plotted parameter space.

For Dirac neutrinos, again inverted hierarchy seems to be ruled out as can be easily checked from figures 2a, 2b and 2c, plotted in a manner similar to the Majorana case

by constraining one mixing angle by its experimental limits and plotting the parameter space for the other two angles. Again, these figures reveal that the plotted parameter space does not overlap with the experimental limits of at least one of the plotted angles, thereby indicating that inverted hierarchy is ruled out at 3σ C.L. for Dirac neutrinos as well.

For Majorana or Dirac neutrinos the case of neutrino masses being degenerate, characterized by either $m_{\nu_1} \lesssim m_{\nu_2} \sim m_{\nu_3} \lesssim 0.1$ eV or $m_{\nu_3} \sim m_{\nu_1} \lesssim m_{\nu_2} \lesssim 0.1$ eV corresponding to normal hierarchy and inverted hierarchy respectively, is again ruled out. Considering degenerate scenario corresponding to inverted hierarchy, figures 1 and 2 can again be used to rule out degenerate scenario at 3σ C.L. for Majorana and Dirac neutrinos respectively. It needs to be mentioned that while plotting these figures the range of the lightest neutrino mass is taken to be 10^{-8} eV – 10^{-1} eV, which includes the neutrino masses corresponding to degenerate scenario, therefore by discussion similar to the one given for ruling out inverted hierarchy, degenerate scenario of neutrino masses is ruled out as well.

Coming to degenerate scenario corresponding to normal hierarchy, one can easily show that this is ruled out again. To this end, in figure 3, by giving full variation to other parameters, we have plotted the mixing angle θ_{12} against the lightest neutrino mass m_{ν_1} . Figure 3a corresponds to the case of Majorana neutrinos and figure 3b to the case of Dirac neutrinos. From the figures one can immediately find that the values of θ_{12} corresponding to $m_{\nu_1} \lesssim 0.1$ eV lie outside the experimentally allowed range, thereby ruling out degenerate scenario for Majorana as well as Dirac neutrinos at 3σ C.L..

The presence of a few isolated points near the experimentally allowed 3σ regions shown in figures 1 and 2 may raise doubts about our conclusions. In order to check whether there are any solution points within the experimentally allowed 3σ region of plotted angles, we have attempted to obtain a common parameter space pertaining to the three mixing angles simultaneously. Interestingly, we find that all possible cases considered here pertaining to inverted hierarchy and degenerate scenario are again ruled out. It may also be added that in the case when charged leptons are in the flavor basis, one can easily check that inverted hierarchy and degenerate scenarios for the texture 4 zero mass matrices are again ruled out, in agreement with the conclusions of [18]. The results pertaining to this case can easily be derived from our earlier cases.

After ruling out the cases pertaining to inverted hierarchy and degenerate scenario, we now discuss the normal hierarchy cases. For the charged lepton mass matrix M_l being Fritzsch-like or in the flavor basis, for Majorana as well as Dirac neutrinos, in table (1) we have presented the viable ranges of neutrino masses, mixing angles θ_{12} , θ_{23} and θ_{13} , Jarlskog's rephasing invariant parameter J , CP violating phase δ and effective neutrino mass $\langle m_{ee} \rangle$. The parameter J can be calculated by using its expression given in [14], whereas δ can be determined from $J = s_{12}s_{23}s_{13}c_{12}c_{23}c_{13}^2 \sin \delta$ where $c_{ij} = \cos \theta_{ij}$ and $s_{ij} = \sin \theta_{ij}$, for $i, j = 1, 2, 3$. The effective Majorana mass, related to neutrinoless double beta decay $(\beta\beta)_{0\nu}$, is given as

$$\langle m_{ee} \rangle = m_{\nu_1} U_{e1}^2 + m_{\nu_2} U_{e2}^2 + m_{\nu_3} U_{e3}^2. \quad (23)$$

Considering the texture 4 zero case when $M_{\nu D}$ and M_l both have parallel texture structures, a close look at table (1) reveals several interesting points. For both Dirac or Majorana neutrinos, the viable range of the lightest neutrino mass m_{ν_1} is quite different, in particular the range corresponding to Dirac neutrinos is much wider at both the ends as compared to the Majorana neutrinos. Similar conclusions can be arrived at by studying the implications of the well known mixing angle θ_{12} on the lightest neutrino mass m_{ν_1} through a closer look at the figures 3a and 3b. Therefore, a measurement of m_{ν_1} could have important implications for the nature of neutrinos. Somewhat constrained range of m_{ν_2} for the Majorana case as compared to the Dirac case is also due to the constrained range of m_{ν_1} for the Majorana case. Also, from the table one finds that the lower limit on θ_{13} for the Dirac case is considerably lower than for the Majorana case, therefore a measurement of θ_{13} would have important implications for this case. It must be noted that the calculated values of $\langle m_{ee} \rangle$ are much less compared to the present limits of $\langle m_{ee} \rangle$ [19], therefore, these do not have any implications for the texture 4 zero cases considered here. However, the future experiments with considerably higher sensitivities, aiming to measure $\langle m_{ee} \rangle \simeq 3.6 \times 10^{-2}$ eV (MOON [20]) and $\langle m_{ee} \rangle \simeq 2.7 \times 10^{-2}$ eV (CUORE [21]), would have implications on the cases considered here. The different cases of Dirac and Majorana neutrinos do not show any divergence for the ranges of Jarlskog's rephasing invariant parameter .

It may be of interest to construct the Pontecorvo-Maki-Nakagawa-Sakata (PMNS) mixing matrix [22] which for Majorana neutrinos is

$$U = \begin{pmatrix} 0.7599 - 0.8701 & 0.4797 - 0.6294 & 0.0199 - 0.1994 \\ 0.1673 - 0.5715 & 0.3948 - 0.7606 & 0.5720 - 0.8224 \\ 0.1854 - 0.5912 & 0.3549 - 0.7363 & 0.5540 - 0.8094 \end{pmatrix}, \quad (24)$$

wherein we have given the magnitude of the matrix elements. Similarly, for Dirac neutrinos, the PMNS matrix is

$$U = \begin{pmatrix} 0.7604 - 0.9213 & 0.3887 - 0.6317 & 0.0015 - 0.1993 \\ 0.1475 - 0.5552 & 0.4049 - 0.8170 & 0.4154 - 0.8244 \\ 0.1830 - 0.6022 & 0.3648 - 0.7441 & 0.5546 - 0.9095 \end{pmatrix}. \quad (25)$$

A general look at the two matrices reveals that the ranges of the matrix elements are more wider in the case of Dirac neutrinos as compared to those in the case of Majorana neutrinos. A comparison of the two matrices shows that the lower limit of the element $U_{\mu 3}$ show an appreciable difference, which seems to be due to the nature of neutrinos, hence, a further precision of $U_{\mu 3}$ would have important implications. Also, it may be mentioned that both the above mentioned matrices are fully compatible with a very recent construction of a mixing matrix by Bjorken *et al.* [23] assuming democratic trimaximally mixed ν_2 mass eigenstate as well as with the one presented by Giunti [24].

For the sake of completion pertaining to normal hierarchy, in table (1) we have also presented the results when charged leptons are in the flavor basis which can be easily deduced from the case when M_l is Fritzsch-like. Interestingly, from the table one im-

diately finds that in this case both J and δ are vanishingly small for the wide range of parameters considered here, which can easily be understood by examining the corresponding mixing matrix. Also, the range of angle θ_{13} is much narrower compared to the case when M_l is Fritzsch-like, particularly for the Majorana case the predicted range is very narrow, therefore a measurement of θ_{13} would have an immediate impact on this case. It may also be added that for the Majorana case, the range of θ_{23} is compatible only with the lower part of the present admissible range, however for the Dirac case there is no such restriction. These conclusions are broadly in agreement with those of [18].

To summarize, detailed calculations have been carried out for different hierarchies in the case of Fritzsch-like texture 4 zero mass matrices with parallel texture structures for charged leptons and for Dirac neutrino mass matrix ($M_{\nu D}$) using latest 3σ input values of neutrino mass and mixing parameters. For the inverted hierarchy, pertaining to both Majorana and Dirac neutrinos, parameter space available to any two of the mixing angles has been explored while considering wide ranges of free parameters available. Similarly, the viability of ‘different’ degenerate scenarios has been examined. Interestingly for both types of neutrinos, inverted hierarchy as well as degenerate scenarios seem to be ruled out at 3σ C.L. and hence strongly disfavored. It may also be added that the results when charged leptons are in the flavor basis can easily be deduced from the present calculations and these lead to similar conclusions.

For the normal hierarchy case, several phenomenological quantities such as Jarlskog’s rephasing invariant parameter J , the CP violating Dirac-like phase δ and the effective neutrino mass $\langle m_{ee} \rangle$ have been calculated. The different cases of Majorana and Dirac neutrinos do not show any divergence for the ranges of J and phase δ . In the case of m_{ν_1} and θ_{13} , the Dirac case seems to accommodate a larger range of these parameters. In particular, a measurement of the lower limits of these parameters would have implications for the nature of neutrinos. Also, the PMNS matrices constructed for Majorana as well as Dirac neutrinos, by giving full variation to the parameters, are compatible with a very recent construction of a mixing matrix by Bjorken *et al.* [23] assuming democratic trimaximally mixed ν_2 mass eigenstate.

Acknowledgements

M.G. and G.A. would like to thank DAE, BRNS (grant No.2005/37/4/BRNS), India, for financial support. S.K. acknowledges the financial support provided by CSIR, India. M.R. would like to thank the Director, UIET for providing facilities to work.

References

- [1] R. Davis, Prog. Part. Nucl. Phys. 32, (1994) 13 ; B. T. Cleveland *et al.*, Astrophys. J. 496, (1998) 505 ; J.N. Abdurashitov *et al.*, SAGE Collaboration, Phys. Rev. C60, (1999) 055801; C. M. Cattadori, GNO Collaboration, Nucl. Phys. Proc. Suppl. 110, (2002) 311 ; W. Hampel *et al.*, GALLEX Collaboration, Phys. Lett. B447, (1999) 127; S. Fukuda *et al.*, Super-Kamiokande Collaboration, Phys. Lett. B539, (2002)

- 179 ; Q. R. Ahmad *et al.*, SNO Collaboration, Phys. Rev. Lett. 89, (2002) 011301; S. N. Ahmad *et al.*, Phys. Rev. Lett. 92, (2004) 181301.
- [2] Y. Fukuda *et al.*, SuperKamiokande Collaboration, Phys. Rev. Lett. 81, (1998) 1562; A. Surdo, MACRO Collaboration, Nucl. Phys. Proc. Suppl. 110, (2002) 342; M. Sanchez, Soudan 2 Collaboration, Phys. Rev. D68, (2003) 113004.
- [3] K. Eguchi *et al.*, KamLAND Collaboration, Phys. Rev. Lett. 90, (2003) 021802; Phys. Rev. Lett. 94, (2005) 081801.
- [4] M. H. Ahn *et al.*, K2K Collaboration, Phys. Rev. Lett. 90, (2003) 041801.
- [5] H. Fritzsch, Z. Z. Xing, Prog. Part. Nucl. Phys. 45, (2000) 1, and references therein.
- [6] H. Fritzsch, Z. Z. Xing, Prog. Part. Nucl. Phys. J. C36, (2004) 227, and references therein.
- [7] K. Kang, S.K. Kang, Phys. Rev. D56, (1997) 1511; H. Nishiura, K. Matsuda, T. Fukuyama, Phys. Rev. D60, (1999) 013006; K. Matsuda, T. Fukuyama, H. Nishiura, Phys. Rev. D61, (2000) 053001; K. Kang, S.K. Kang, C.S. Kim, S.M. Kim, Mod. Phys. Lett. A16, (2001) 2169; C. Giunti, M. Tanimoto, Phys. Rev. D66, (2002) 113006; M. Frigerio, A.Yu. Smirnov, Nucl. Phys. B640, (2002) 233; P.F. Harrison, W.G. Scott, Phys. Lett. B547, (2002) 219; E. Ma, Mod. Phys. Lett. A17, (2002) 2361; E. Ma, Phys. Rev. D66, (2002) 117301; P.H. Frampton, S.L. Glashow, D. Marfatia, Phys. Lett. B 536, (2002) 79; Z.Z. Xing, Phys. Lett. B 530, (2002) 159; Z.Z. Xing, Phys. Lett. B 539, (2002) 85; A. Kageyama, S. Kaneko, N. Simoyama, M. Tanimoto, Phys. Lett. B 538, (2002) 96; W.L. Guo, Z.Z. Xing, hep-ph/0211315; M. Randhawa, G. Ahuja, M. Gupta, Phys. Rev. D65, (2002) 093016; M. Frigerio, A.Yu. Smirnov, Phys. Rev. D67, (2003) 013007; K.S. Babu, E. Ma, J.W.F. Valle, Phys. Lett. B552, (2003) 207; W.L. Guo, Z.Z. Xing, Phys. Rev. D67, (2003) 053002; S. Kaneko, M. Tanimoto, Phys. Lett. B551, (2003) 127; B.R. Desai, D.P. Roy, A.R. Vaucher, Mod.Phys.Lett. A18, (2003) 1355; K. Hasegawa, C.S. Lim, K. Ogure, Phys. Rev. D68, (2003) 053006; M. Honda, S. Kaneko, M. Tanimoto, JHEP 0309, (2003) 028; G. Bhattacharyya, A. Raychaudhuri, A. Sil, Phys. Rev. D67, (2003) 073004; P.F. Harrison, W.G. Scott, Phys. Lett. B594, (2004) 324; Z.Z. Xing, Int. J. Mod. Phys. A19, (2004) 1; M. Bando, S. Kaneko, M. Obara, M. Tanimoto, Phys. Lett. B580, (2004) 229; O.L.G. Peres, A.Yu. Smirnov, Nucl. Phys. B680, (2004) 479; C.H. Albright, Phys. Lett. B599, (2004) 285; J. Ferrandis, S. Pakvasa, Phys. Lett. B603, (2004) 184; S.T. Petcov, W. Rodejohann, Phys. Rev. D71, (2005) 073002; S.S. Ma-sood, S. Nasri, J. Schechter, Phys. Rev. D71, (2005) 093005; R. Dermíšek, S. Raby, Phys. Lett. B622, (2005) 327 ; F. Plentinger, W. Rodejohann, Phys. Lett. B625, (2005) 264.
- [8] P. Minkowski, Phys. Lett. B67, (1977) 421; T. Yanagida, in Proc. of Work-shop on Unified Theory and Baryon number in the Universe, eds. O. Sawada, A. Sugamoto,

- KEK, Tsukuba, (1979); M. Gell-Mann, P. Ramond, R. Slansky, in Supergravity, eds P. van Nieuwenhuizen, D. Z. Freedman (North Holland, Amsterdam 1980); P. Ramond, Sanibel talk, retroprinted as hep-ph/9809459; S. L. Glashow, in Quarks and Leptons, Cargèse lectures, eds M. Levy, (Plenum, 1980, New York) p. 707; R. N. Mohapatra, G. Senjanovic, Phys. Rev. Lett. 44, (1980) 912.
- [9] Z. Z. Xing, H. Zhang, Phys.Lett. B569, (2003) 30; Z. Z. Xing, S. Zhou, Phys. Lett. B606, (2005) 145.
- [10] K. Matsuda, H. Nishiura, Phys. Rev. D74, (2006) 033014, and references therein.
- [11] D. Du, Z. Z. Xing, Phys. Rev. D48, (1993) 2349; L.J. Hall, A. Rasin, Phys. Lett. B315, (1993) 164 ; P. Ramond, R. G. Roberts, G. G. Ross, Nucl. Phys. B406, (1993) 19; H. Fritzsch, D. Holtmanspötter, Phys. Lett. B338, (1994) 290; H. Fritzsch, Z.Z. Xing, Phys. Lett. B353, (1995) 114; M. Gupta, P. S. Gill, Pramana 45, (1995) 333; P.S. Gill, M. Gupta, J. Phys. G: Nuc. Part. Phys. 21, (1995) 1; H. Lehmann, C. Newton, T. T. Wu, Phys. Lett. B384, (1996) 249; P. S. Gill, M. Gupta, Jol. Phys. G23, (1997) 1; P. S. Gill, M. Gupta, Phys. Rev. D56, (1997) 3143; Z. Z. Xing, J. Phys. G23, (1997) 1563; T. Kobayashi, Z. Z. Xing, Mod. Phys. Lett. A12, (1997) 561; *ibid.* Int. J. Mod. Phys. A13, (1998) 2201; P. S. Gill, M. Gupta, Phys. Rev. D57, (1998) 3917; J. L. Chkareuli, C. D. Froggatt, Phys. Lett. B450, (1999) 158; A. Mondragón, E. Rodriguez-Jáuregui, Phys. Rev. D59, (1999) 093009; M. Randhawa, V. Bhatnagar, P. S. Gill, M. Gupta, Phys. Rev. D60, (1999) 051301; G. C. Branco, D. Emmanuel-Costa, R. González Felipe, Phys. Lett. B477, (2000) 147; S. H. Chiu, T. K. Kuo, G. H. Wu, Phys. Rev. D62, (2000) 053014; H. Fritzsch, Z. Z. Xing, Phys. Rev. D61, (2000) 073016; *ibid.* Phys. Lett. B506, (2001) 109; R. Rosenfeld, J. L. Rosner, Phys. Lett. B516, (2001) 408 ; R. G. Roberts, A. Romanino, G. G. Ross, L. Velasco-Sevilla, Nucl. Phys. B615, (2001) 358; J. L. Chkareuli, C. D. Froggatt, Nucl. Phys. B626, 307 (2002); H. Fritzsch, Z. Z. Xing, Phys. Lett. B555, (2003) 63; Z. Z. Xing, H. Zhang, J. Phys. G30, (2004) 129.
- [12] W. Grimus, hep-ph/0511078.
- [13] See M. Randhawa *et al.* from [11].
- [14] M. Fukugita, M. Tanimoto, T. Yanagida, Prog. Theor. Phys. 89, (1993) 263; *ibid.* Phys. Lett. B562, (2003) 273.
- [15] J. W. F. Valle, hep-ph/0608101; *ibid.* hep-ph/0612251.
- [16] M.C. Gonzalez-Garcia, M. Maltoni, arXiv:0704.1800v1.
- [17] M. Appolonio *et al.*, CHOOZ Collaboration, Eur. Phys. J. C27 (2003) 331.
- [18] S. Dev, S. Kumar, S. Verma, S. Gupta, hep-ph/0612102.

- [19] H. V. Klapdor-Kleingrothaus *et al.*, Heidelberg-Moscow collaboration, Eur. Phys. J. A12, (2001) 147; *ibid.* Phys. Lett. B586, (2004) 198, and references therein.
- [20] H. Ejiri, R. Hazama, P. Krastev, N. Kudomi, R. G. H. Robertson, Phys. Rev. Lett. 85, (2000) 2917.
- [21] R. Ardito, *et al.*, CUORE Collaboration, hep-ex/0501010; C. Arnaboldi *et al.*, Phys. Rev. Lett. 95, (2005) 142501.
- [22] B. Pontecorvo, Zh. Eksp. Teor. Fiz. (JETP) 33, (1957) 549; *ibid.* 34, (1958) 247; *ibid.* 53, (1967) 1771; Z. Maki, M. Nakagawa, S. Sakata, Prog. Theor. Phys. 28, (1962) 870.
- [23] J. D. Bjorken, P. F. Harrison, W. G. Scott, Phys. Rev. D74, (2006) 073012.
- [24] C. Giunti, hep-ph/0611125.

	M_l being Fritzsch-like		M_l being in the flavor basis	
	Dirac case	Majorana case	Dirac case	Majorana case
m_{ν_1}	$5.73 \times 10^{-5} - 0.012$	$2.47 \times 10^{-4} - 0.006$	$(1.63 - 6.28) \times 10^{-3}$	$(.402 - 2.06) \times 10^{-3}$
m_{ν_2}	0.0084 - 0.0149	0.0084 - 0.0108	0.0086 - 0.0113	0.0084 - 0.0096
m_{ν_3}	0.0456 - 0.0577	0.0455 - 0.0575	0.0446 - 0.0576	0.0455 - 0.0573
θ_{12}	$29.30^\circ - 39.20^\circ$	$29.30^\circ - 39.20^\circ$	$29.30^\circ - 39.20^\circ$	$29.30^\circ - 39.04^\circ$
θ_{23}	$35.70^\circ - 55.60^\circ$	$35.70^\circ - 55.60^\circ$	$35.70^\circ - 55.59^\circ$	$35.70^\circ - 40.15^\circ$
θ_{13}	$0.084^\circ - 11.50^\circ$	$1.14^\circ - 11.50^\circ$	$3.60^\circ - 11.15^\circ$	$8.43^\circ - 11.50^\circ$
J	-0.0462 - .0448	-0.0459 - .0463	~ 0	~ 0
δ	$-90^\circ - 90.0^\circ$	$-90^\circ - 90.0^\circ$	$\sim 0^\circ$	$\sim 0^\circ$
$\langle m_{ee} \rangle$	-	.00086 - .0173	-	.0032 - .0075

Table 1: Calculated ranges for neutrino mass and mixing parameters obtained by varying ϕ_1 and ϕ_2 from 0 to 2π for the normal hierarchy case. Inputs have been defined in the text. All masses are in eV.




Figure 1: Plots showing the parameter space corresponding to any of the two mixing angles by constraining the third angle by its experimental limits given in equation (22) and giving full allowed variation to other parameters for Majorana neutrinos. The blank rectangular region indicates the experimentally allowed 3σ region of the plotted angles.




Figure 2: Plots showing the parameter space corresponding to any of the two mixing angles by constraining the third angle by its experimental limits given in equation (22) and giving full allowed variation to other parameters for Dirac neutrinos. The blank rectangular region indicates the experimentally allowed 3σ region of the plotted angles.




Figure 3: Plots showing variation of mixing angle θ_{12} with lightest neutrino mass m_{ν_1} by giving full variation to other parameters for neutrinos being (a) Majorana-like and (b) Dirac-like. The parallel lines indicate the 3σ limits of angle θ_{12} .